

DISULFIRAM

Patient Treatment Card



Patient Information

You have been prescribed Disulfiram tablets and you must not consume any alcohol as long as you are taking this medicine and for at least 14 days after stopping treatment. If you are taking other medicines particularly cough syrups and tonics you should check with your doctor or pharmacist to make sure they are suitable to take with Disulfiram.

Certain foods, liquid medicines, remedies, tonics, toiletries, perfumes and sprays may contain enough alcohol to cause a Disulfiram-alcohol reaction. Caution should also be exercised with alcohol-free or low-alcohol beers and wines which may provoke a reaction when consumed in sufficient quantities. Please refer to patient information leaflet for details.

Please carry this card with you at all times to make sure you receive the correct treatment in the event of an accident or sudden illness.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard

By reporting side effects, you can help provide more information on the safety of this medicine. Adverse events may also be reported directly to Teva UK Limited at www.tevauk.com

Healthcare Professional Information

This patient should not be given alcohol in any form until 14 days after stopping Disulfiram therapy. Disulfiram interferes with drug metabolizing enzymes and may potentiate the action of centrally acting drugs, (see SPC for full details).

Disulfiram therapy patient Details

Daily Dose

Name

Address

Telephone (day)

(evening)

Date of issue

Doctor's Name

Doctors telephone

October 2019

UK/GEN/16/0056(2)