

ROSUVASTATIN 5 mg, 10 mg, 20 mg AND 40 mg FILM-COATED TABLETS

Package leaflet: Information for the patient

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist. This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Rosuvastatin is and what it is used for
2. What you need to know before you take Rosuvastatin
3. How to take Rosuvastatin
4. Possible side effects
5. How to store Rosuvastatin
6. Contents of the pack and other information

1 What Rosuvastatin is and what it is used for

Rosuvastatin belongs to a group of medicines called statins.

You have been prescribed Rosuvastatin because:

- you have a high cholesterol level. This means you are at risk from a heart attack or stroke. Rosuvastatin is used in adults, adolescents and children 8 years or older to treat high cholesterol. You have been advised to take a statin, because changing your diet and doing more exercise were not enough to correct your cholesterol levels. You should continue with your cholesterol-lowering diet and exercise while you are taking Rosuvastatin.

- Or you have other factors that increase your risk of having a heart attack, stroke or related health problems.

Heart attack, stroke and other problems can be caused by a disease called atherosclerosis. Atherosclerosis is due to build up of fatty deposits in your arteries.

Why is it important to keep taking Rosuvastatin?

Rosuvastatin is used to correct the levels of fatty substances in the blood called lipids, the most common of which is cholesterol.

- There are different types of cholesterol found in the blood – 'bad' cholesterol (LDL-C) and 'good' cholesterol (HDL-C).
- Rosuvastatin can reduce the 'bad' cholesterol and increase the 'good' cholesterol.
- It works by helping to block your body's production of 'bad' cholesterol. This means your body's ability to remove it from your blood.

For most people, high cholesterol does not affect their health, because it does not produce any symptoms.

However, if it is left untreated, fatty deposits can build up in the walls of your blood vessels causing them to narrow. Sometimes, these narrowed blood vessels can get blocked which can cut off the blood supply to the heart or brain leading to a heart attack or a stroke. By lowering your cholesterol levels, you can reduce your risk of having a heart attack, a stroke or related health problems.

You need to keep taking

Rosuvastatin, even if it has got your cholesterol to the right level, because it prevents your cholesterol levels from creeping up again and causing build up of fatty deposits. However, you should stop if your doctor tells you to do so, or you have become pregnant.

2 What you need to know before you take Rosuvastatin

Do not take Rosuvastatin if you:

- are allergic to rosuvastatin or any of the other ingredients of this medicine (listed in section 6)
- have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking rosuvastatin or other related medicines.
- are pregnant or breast-feeding. If you become pregnant while taking Rosuvastatin stop taking it immediately and tell your doctor. Women should avoid becoming pregnant while taking Rosuvastatin by using suitable contraception.
- have liver disease
- have severe kidney problems
- have repeated or unexplained muscle aches or pains
- take a drug combination of sofosbuvir/velpatasvir/voxilaprevir (used for viral infection of the liver called hepatitis C)
- take a drug called ciclosporin (used, for example, after organ transplants)

If any of the above applies to you (or you are in doubt), please go back and see your doctor.

In addition, do not take Rosuvastatin 40 mg Film-coated Tablets (the highest dose) if:

- you have moderate kidney problems (if in doubt, please ask your doctor)
- your thyroid gland is not working properly
- you have had any repeated or unexplained muscle aches or pains, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines
- you regularly drink large amounts of alcohol
- you are of Asian origin (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian)
- you take other medicines called fibrates to lower your cholesterol.

If any of the above applies to you (or you are in doubt), please go back and see your doctor.

Warnings and precautions

Talk to your doctor or pharmacist before taking Rosuvastatin if:

- you have problems with your kidneys
- you have problems with your liver
- you have had repeated or unexplained muscle aches or pains, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines. Tell your doctor immediately if you have unexplained muscle aches or pains especially if you feel unwell or have a fever. Also tell your doctor or pharmacist if you have a muscle weakness that is constant.

- If you have or have had myasthenia (a disease with general muscle weakness including in some cases muscles used when breathing), or ocular myasthenia (a disease causing eye muscle weakness) as statins may sometimes aggravate the condition or lead to the occurrence of myasthenia (see section 4).
- you regularly drink large amounts of alcohol
- your thyroid gland is not working properly
- you take other medicines called fibrates to lower your cholesterol. Please read this leaflet carefully, even if you have taken other medicines for high cholesterol before you take medicines used to treat the HIV infection e.g. ritonavir with lopinavir and/or atazanavir, please see Other medicines and Rosuvastatin

- you are taking or have taken in the last 7 days a medicine called fusidic acid, (a medicine for bacterial infection) orally or by injection. The combination of fusidic acid and Rosuvastatin can lead to serious muscle problems (rhabdomyolysis), please see Other medicines and Rosuvastatin film-coated tablets.
- you are over 70 (your doctor needs to choose the right start dose of Rosuvastatin to suit you).
- you have severe respiratory failure.
- you are of Asian origin - that is Japanese, Chinese, Filipino, Vietnamese, Korean and Indian. Your doctor needs to choose the right start dose of Rosuvastatin to suit you.

If any of the above applies to you (or if you are not sure):

- Do not take Rosuvastatin 40 mg Film-coated Tablets (the highest dose) and check with your doctor or pharmacist before you actually start taking any dose of Rosuvastatin Film-coated Tablets.

In a small number of people, statins can affect the liver. This is identified by a simple test which looks at increased levels of liver enzymes in the blood. For this reason, your doctor will usually carry out this blood test (liver function test) before and during treatment with Rosuvastatin Film-coated Tablets.

Serious skin reactions including Stevens-Johnson syndrome and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with rosuvastatin treatment. Stop using Rosuvastatin Film-coated tablets and seek medical attention immediately if you notice any of the symptoms described in section 4.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

Children and adolescents

- If the patient is under 6 years old: Rosuvastatin Film-coated tablets should not be given to children younger than 6 years.
- If the patient is below 18 years of age: The Rosuvastatin 40 mg tablet is not suitable for use in children and adolescents below 18 years of age.

Other medicines and Rosuvastatin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor if you are taking any of the following:

- ciclosporin (used for example, after organ transplants)
- warfarin, clopidogrel, ticagrelor (or any other drug used for thinning the blood)

- fibrates (such as gemfibrozil, fenofibrate) or other medicine used to lower cholesterol (such as ezetimibe)
- indigestion remedies (used to neutralise acid in your stomach)
- erythromycin (an antibiotic)
- fusidic acid (an antibiotic). If you need to take oral fusidic acid to treat bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Rosuvastatin. Taking Rosuvastatin with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in section 4.

- an oral contraceptive (the pill)
- regorafenib (used to treat cancer)
- darolutamide (used to treat cancer),
- capmatinib (used to treat cancer)
- hormone replacement therapy
- fostamatinib (used to treat low platelet counts)
- febuxostat (used to treat and prevent high blood levels of uric acid)
- teriflunomide (used to treat multiple sclerosis)

- any of the following drugs used to treat viral infections, including HIV or hepatitis C infection, alone or in combination (please see Warnings and precautions): ritonavir, lopinavir, atazanavir, sofosbuvir, voxilaprevir, ombitasvir, paritaprevir, dasabuvir, velpatasvir, grazoprevir, elbasvir, glecaprevir, pibrentasvir.

The effects of these medicines could be changed by Rosuvastatin or they could change the effect of Rosuvastatin.

Pregnancy and breast-feeding

Do not take Rosuvastatin if you are pregnant or breast-feeding. If you become pregnant while taking Rosuvastatin stop taking it immediately and tell your doctor. Women should avoid becoming pregnant while taking Rosuvastatin by using suitable contraception.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Most people can drive a car and operate machinery while using Rosuvastatin – it will not affect their ability. However, some people feel dizzy during treatment with Rosuvastatin. If you feel dizzy, stop your doctor before attempting to drive or use machines.

Rosuvastatin Film-coated Tablets contains lactose and sodium.

Lactose If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Sodium This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3 How to take Rosuvastatin

Always take this medicine as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Usual dose in adults

If you are taking Rosuvastatin for high cholesterol:

Starting dose Your treatment with Rosuvastatin must start with the **5 mg or the 10 mg dose**, even if you have taken a higher dose of a different statin before. The choice of your start dose will depend upon:

- your cholesterol levels
 - the level of risk you have of experiencing a heart attack or stroke
 - whether you have a factor that may make you more sensitive to possible side effects.
- Please check with your doctor or pharmacist which start dose of Rosuvastatin will best suit you.

Your doctor may decide to give you the lowest dose (5 mg) if you:

- you are of Asian origin (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian)
- are over 70 years of age
- have moderate kidney problems
- are at risk of muscle aches and pains (myopathy).

Increasing the dose and maximum daily dose

Your doctor may decide to increase your dose. This is so that you are taking the amount of Rosuvastatin that is right for you. If you started with a 5 mg dose, your doctor may decide to double this to 10 mg, then 20 mg and then 40 mg if necessary. If you started on 10 mg, your doctor may decide to double this to 20 mg and then 40 mg if necessary. There will be a gap of four weeks between every dose adjustment. The maximum daily dose of Rosuvastatin Film-coated Tablets is 40 mg. It is only for patients with high cholesterol levels and a high risk of heart attacks or stroke, whose cholesterol levels are not lowered enough with 20 mg.

If you are taking Rosuvastatin to reduce your risk of having a heart attack, stroke or related health problems:

The recommended dose is 20 mg daily. However, your doctor may decide to use a lower dose if you have any of the factors mentioned above.

Use in children and adolescents aged 6-17 years

The dose range in children and adolescents aged 6 to 17 years is 5 to 20 mg once daily. The usual start dose is 5 mg per day and your doctor may gradually increase your dose to find the right amount of Rosuvastatin Film-coated Tablets for you. The maximum daily dose of Rosuvastatin Film-coated Tablets is 10 mg or 20 mg depending on your age and depending on your underlying condition being treated. Take your dose once a day. Rosuvastatin 40 mg Film-coated Tablets should not be used by children.

Taking your tablets

Swallow each tablet whole with a drink of water.

Take Rosuvastatin Film-coated Tablets once daily.

You should take your dose at any time of the day with or without food. Try to take your tablet at the same time every day to help you to remember it.

Regular cholesterol checks

It is important to go back to your doctor for regular cholesterol checks to make sure your cholesterol has reached and is staying at the correct level.

Your doctor may decide to increase your dose so that you are taking the amount of Rosuvastatin Film-coated Tablets that is right for you.

If you take more Rosuvastatin than you should

Contact your doctor or nearest hospital for advice.

If you go into hospital or receive treatment for another condition, tell the medical staff that you are taking Rosuvastatin Film-coated Tablets.

If you forget to take Rosuvastatin Film-coated Tablets

Do not worry, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Rosuvastatin

Stopping Tablets If you want to stop taking Rosuvastatin. Your cholesterol levels might increase again if you stop taking Rosuvastatin. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

It is important that you are aware of what side effects may be. They are usually mild and disappear after a short time.

Stop taking Rosuvastatin and seek medical help immediately if you have any of the following allergic reactions:

- difficulty in breathing, with or without swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing
- severe itching of the skin (with raised lumps).
- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by flu-like and flu-like symptoms (Stevens-Johnson syndrome)
- widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

Also, stop taking Rosuvastatin and talk to your doctor immediately:

- if you have any unusual aches or pains in your muscles which go on for longer than you might expect. Muscle symptoms are more common in children and adolescents than in adults. As with other statins, a very small number of people have experienced severe muscle aches and pains and rarely these have gone on to become a potentially life threatening muscle damage known as rhabdomyolysis.
- if you experience muscle rupture
- if you have lupus-like disease syndrome (including rash, joint problems and effects on blood cells)

Common: possible side effects (may affect up to 1 in 10 people):

- headache
- stomach pain
- constipation
- feeling sick
- muscle pain
- feeling weak
- dizziness
- an increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking Rosuvastatin (only Rosuvastatin 40 mg Film-coated Tablets)
- diabetes. This is more likely if you have high levels of sugars and fat in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.

Uncommon possible side effects (may affect up to 1 in 100 people):

- rash, itching or other skin reactions
- an increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking Rosuvastatin (only Rosuvastatin 5 mg, 10 mg and 20 mg Film-coated Tablets)

Rare possible side effects (may affect up to 1 in 1,000 people):

- severe allergic reaction – signs include swelling of the face, lips, tongue and/or throat, difficulty in swallowing and breathing, a severe itching of the skin (with raised lumps). If you think you are having an allergic reaction, then stop taking Rosuvastatin and seek medical help immediately
- muscle damage in adults – as a precaution, stop taking Rosuvastatin and talk to your doctor immediately if you have any unusual aches or pains in your muscles which go on for longer than expected
- a severe stomach pain (inflamed pancreas)
- increase in liver enzymes in the blood
- bleeding or bruising more easily than normal due to low level of blood platelets
- lupus-like disease syndrome (including rash, joint disorders and effects on blood cells)

Very rare possible side effects (may affect up to 1 in 10,000 people):

- jaundice (yellowing of the skin and eyes)
- hepatitis (an inflamed liver)
- traces of blood in your urine
- damage to the nerves of your legs and arms (such as numbness)
- cold pain
- memory loss
- breast enlargement in men (gynaecomastia).

Side effects of unknown frequency may include:

- diarrhoea (loose stools)
 - cough
 - shortness of breath
 - oedema (swelling)
 - sleep disturbances including insomnia and nightmares
 - sexual difficulties
 - depression
 - breathing problems including persistent cough and/or shortness of breath or fever
 - tendon injury
 - muscle weakness that is constant.
 - myasthenia gravis (a disease causing general muscle weakness including some cases muscles used when breathing)
 - ocular myasthenia (a disease causing eye muscle weakness)
- Talk to your doctor if you experience weakness in your arms or legs that worsens after periods of activity, double vision or drooping of your eyelids, difficulty swallowing, or shortness of breath.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Rosuvastatin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Blisters: Do not store above 30°C.

Bottles: This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Rosuvastatin Film-coated Tablets contains

The active substance is rosuvastatin.

Rosuvastatin 5 mg, 10 mg, 20 mg & 40 mg Film-coated Tablets contain rosuvastatin calcium equivalent to 5 mg, 10 mg, 20 mg or 40 mg of rosuvastatin.

The other ingredients are:

Tablet core: microcrystalline cellulose, lactose monohydrate, croscopolone (type B), hydroxypropylcellulose, sodium hydrogencarbonate & magnesium stearate.

Film-coating: lactose monohydrate, hypromellose 6 Cp, titanium dioxide (E171), triacetin, iron oxide yellow (E172) (Rosuvastatin 5 mg Film-coated Tablets) & iron oxide red (E172) (Rosuvastatin 10 mg, 20 mg and 40 mg Film-coated Tablets).

What Rosuvastatin Film-coated Tablets looks like and contents of the pack

Rosuvastatin 5 mg Film-coated Tablets: round, biconvex, yellow tablet, debossed with '5' on one side, around 5.5 mm in diameter.

Rosuvastatin 10 mg Film-coated Tablets: round, biconvex, pink tablet, debossed with '10' on one side and with a score line on the other side, around 7 mm in diameter.

Rosuvastatin 20 mg Film-coated Tablets: round, biconvex, pink tablet, debossed with '20' on one side and with a score line on the other side, around 9 mm in diameter.

Rosuvastatin 40 mg Film-coated Tablets: oval, biconvex, pink tablet, debossed with '40' on one side and with a score line on the other side, dimensions of around 11.5 x 7 mm.

Rosuvastatin 10, 20, 40 mg Film-coated Tablets: The tablets can be divided into equal halves.

Rosuvastatin Film-coated Tablets come in blister packs containing 7, 14, 15, 20, 28, 28 (calendar pack), 30, 30 x 1, 42, 50, 50x1, 56, 60, 84, 90, 90x1, 98, 100 and 100x1 film-coated tablets and HDPE bottles containing 28, 30, 100 and 250 film-coated tablets. Not all pack sizes may be marketed.

Marketing Authorisation Holder Teva UK Limited, Ridings Whistler Drive, Castleford, WF10 5HX, United Kingdom

Manufacturer TEVA PHARMA S.L.U., C/C, nº 4, Zona Industrial, Polígono Industrial Zaragoza, Spain

This leaflet was last revised in October 2023.

PL 00289/1854-1857