

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

GoResp Digihaler 320 micrograms/9 micrograms inhalation powder

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each delivered dose (the dose that leaves the mouthpiece of the Digihaler) contains 320 micrograms of budesonide and 9 micrograms of formoterol fumarate dihydrate.

This is equivalent to a metered dose of 400 micrograms budesonide and 12 micrograms of formoterol fumarate dihydrate.

Excipient(s) with known effect:

Each dose contains approximately 10 milligrams of lactose (as monohydrate).

For the full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

Inhalation powder.

White powder.

White inhaler with a semi-transparent wine red mouthpiece cover and an integrated electronic Module (eModule) for recording, storing and communication (via Bluetooth® technology with a mobile application and a dashboard program, upon patient/user consent) of inhaler usage data.

### **4 CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

GoResp Digihaler is indicated in adults 18 years of age and older only.

##### Asthma

GoResp Digihaler is indicated in the regular treatment of asthma, where use of a combination (inhaled corticosteroid and long-acting  $\beta_2$  adrenoceptor agonist) is appropriate:

-in patients not adequately controlled with inhaled corticosteroids and “as needed” inhaled short-acting  $\beta_2$  adrenoceptor agonists.

or

-in patients already adequately controlled on both inhaled corticosteroids and long-acting  $\beta_2$  adrenoceptor agonists.

##### COPD

Symptomatic treatment of patients with COPD with forced expiratory volume in 1 second (FEV<sub>1</sub>) < 70% predicted normal (post bronchodilator) and a history of repeated exacerbations, who have significant symptoms despite regular therapy with long-acting bronchodilators.

## **4.2 Posology and method of administration**

### Posology

#### Asthma

GoResp Digihaler is not intended for the initial management of asthma.

GoResp Digihaler is not an appropriate treatment for the adult patient with only mild asthma.

The dosage of GoResp Digihaler is individual and should be adjusted to the severity of the disease. This should be considered not only when treatment with combination medicinal products is initiated but also when the maintenance dose is adjusted. If an individual patient should require a combination of doses other than those available in the combination inhaler, appropriate doses of  $\beta_2$  adrenoceptor agonists and/or corticosteroids by individual inhalers should be prescribed.

Once asthma symptoms are controlled, consideration may be given to gradually reducing the dose of GoResp Digihaler. Patients should be reassessed regularly by their prescriber/health care provider so that the dose of GoResp Digihaler remains optimal. The dose should be titrated to the lowest dose at which effective control of symptoms is maintained.

When it is appropriate to titrate down to a lower strength than is available for GoResp Digihaler, a change to an alternative fixed-dose combination of budesonide and formoterol fumarate containing a lower dose of the inhaled corticosteroid is required. When long-term control of symptoms is maintained with the lowest recommended dose, then the next step could include a test of inhaled corticosteroid alone.

In usual practice when control of symptoms is achieved with the twice daily dose regimen with a lower strength product, titration to a lower effective dose could include once daily dosing when, in the opinion of the prescriber, a long-acting bronchodilator is required to maintain control rather than treatment with an inhaled corticosteroid alone.

Patients should be advised to have their separate rapid-acting bronchodilator reliever inhaler available for rescue use at all times.

#### *Recommended doses:*

Adults (18 years and older): 1 inhalation twice daily. Some patients may require up to a maximum of 2 inhalations twice daily.

Increasing use of a separate rapid-acting bronchodilator indicates a worsening of the underlying condition and warrants a reassessment of the asthma therapy.

GoResp Digihaler 320 micrograms/9 micrograms should be used as maintenance therapy only. A lower strength of GoResp Digihaler is available for the maintenance and reliever therapy regimen.

### COPD

### *Recommended doses:*

Adults (18 years and older):

1 inhalation twice daily

### Special populations:

#### *Elderly patients ( $\geq 65$ years old)*

There are no special dosing requirements for elderly patients.

#### *Patients with renal or hepatic impairment*

There are no data available for use of a fixed-dose combination of budesonide and formoterol fumarate dihydrate in patients with hepatic or renal impairment. As budesonide and formoterol are primarily eliminated via hepatic metabolism, an increased exposure can be expected in patients with severe liver cirrhosis.

#### Paediatric population

The safety and efficacy of GoResp Digihaler in children, 12 years and younger and adolescents, 13 to 17 years of age has not yet been established. Currently available data are described in sections 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

This medicinal product is not recommended for use in children and adolescents under the age of 18 years.

### Method of administration

Inhalation use.

*Digihaler* is a breath actuated, inspiratory flow-driven inhaler, with an integrated eModule (a digital version of the commercially available multi-dose dry powder inhaler, *Spiromax*). This means the active substances are delivered into the airways when the patient inhales through the mouthpiece.

The integrated eModule does not impact the metering or dosing functionality of the inhaler. Moderate and severe asthmatic patients were shown to be able to generate sufficient inspiratory flow rate to deliver the therapeutic dose (see section 5.1).

GoResp Digihaler should be used correctly in order to achieve effective treatment. As such, the patients should be advised to read the patient information leaflet carefully and follow the instructions for use as detailed in the leaflet.

The use of GoResp Digihaler follows three steps: open, breathe and close which are outlined below.

**Open:** Hold the Digihaler with the mouthpiece cover at the bottom and open the mouthpiece cover by folding it down until it is fully opened when one click is heard.

**Breathe:** Place the mouthpiece between the teeth with the lips closed around the mouthpiece, do not bite the mouthpiece of the inhaler. Breathe in forcefully and deeply through the mouthpiece. Remove the Digihaler from mouth and hold the breath for 10 seconds or as long as comfortable for the patients.

**Close:** Breathe out gently and close the mouthpiece cover

The Digihaler can be paired with the Digihaler mobile application (App) by scanning the Quick Response code (QR code), located on the top of the inhaler, with a mobile device. The inhaler will still provide the same dose, even if the QR code is not scanned and the inhaler is not connected to the App.

Patients should be advised to use GoResp Digihaler within one minute of opening the wine red mouthpiece cover in order for the data to be recorded electronically. After one minute, the patient can still take their medicine but the data recorded in the App will show low/no inhalation, even if they have performed a good or fair inhalation.

It is important to advise patients to close the mouthpiece cover after taking an inhalation.

It is also important to advise patients not to shake the inhaler before use and not to breathe out through the Digihaler and not to block the air vents when they are preparing the “Breathe” step.

Patients should also be advised to rinse their mouth with water after inhaling (see section 4.4)

The patient may notice a taste when using GoResp Digihaler due to the lactose excipient.

Patients should be advised to not place the inhaler in water.

### **4.3 Contraindications**

Hypersensitivity to the active substances or the excipient listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### General

It is recommended that the dose is tapered when the treatment is discontinued and should not be stopped abruptly.

If patients find the treatment ineffective, or exceed the highest recommended dose of GoResp Digihaler, medical attention must be sought (see section 4.2). Sudden and progressive deterioration in control of asthma or COPD is potentially life-threatening and the patient should undergo urgent medical assessment. In this situation, consideration should be given to the need for increased therapy with corticosteroids, e.g. a course of oral corticosteroids, or antibiotic treatment if an infection is present.

Patients should be advised to have their rescue inhaler available at all times, either GoResp Digihaler (for asthma patients using GoResp Digihaler as maintenance and reliever therapy) or a separate rapid-acting bronchodilator (for asthma patients using GoResp Digihaler as maintenance therapy only).

Patients should be reminded to take their GoResp Digihaler maintenance dose as prescribed, even when asymptomatic. The prophylactic use of GoResp Digihaler, e.g. before exercise, has not been studied. The reliever inhalations of GoResp Digihaler should be taken in response to symptoms but are not intended for regular prophylactic use, e.g. before exercise. For such, a separate rapid-acting bronchodilator should be considered.

### Asthma symptoms

Patients should be reassessed regularly by their prescriber/healthcare provider so that the dose of GoResp Digihaler remains optimal. The dose should be titrated to the lowest dose at which effective control of symptoms is maintained. Once asthma symptoms are controlled, consideration may be given to gradually reducing the dose of GoResp Digihaler. When it is appropriate to titrate down to a lower strength than is available for GoResp Digihaler, a change to an alternative fixed-dose combination of budesonide and formoterol fumarate containing a lower dose of the inhaled corticosteroid is required.

Regular review of patients as treatment is stepped down is important.

Patients should not be initiated on GoResp Digihaler during an exacerbation, or if they have significantly worsening or acutely deteriorating asthma.

Serious asthma-related adverse reactions and exacerbations may occur during treatment with GoResp Digihaler. Patients should be asked to continue treatment but to seek medical advice if asthma symptoms remain uncontrolled or worsen after initiation with GoResp Digihaler.

There are no clinical study data on GoResp Digihaler available in COPD patients with a pre-bronchodilator FEV<sub>1</sub> >50% predicted normal and with a post-bronchodilator FEV<sub>1</sub> <70% predicted normal (see section 5.1)

Paradoxical bronchospasm may occur, with an immediate increase in wheezing and shortness of breath, after dosing. If the patient experiences paradoxical bronchospasm GoResp Digihaler should be discontinued immediately, the patient should be assessed and an alternative therapy instituted, if necessary. Paradoxical bronchospasm responds to a rapid-acting inhaled bronchodilator and should be treated straightaway (see section 4.8).

### Systemic effects

Systemic effects may occur with any inhaled corticosteroid, particularly at high doses prescribed for long periods. These effects are much less likely to occur with inhalation treatment than with oral corticosteroids.

Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma and more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression (particularly in children) (see section 4.8).

#### *Visual disturbance*

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

### Effects on bone density

Potential effects on bone density should be considered, particularly in patients on high doses for prolonged periods that have co-existing risk factors for osteoporosis.

Long-term studies with inhaled budesonide in adults at daily doses of 800 micrograms (metered dose) have not shown any significant effects on bone mineral density. No information regarding the effect of a budesonide/formoterol fumarate dihydrate fixed-dose combination at higher doses is available.

### Adrenal function

If there is any reason to suppose that adrenal function is impaired from previous systemic steroid therapy, care should be taken when transferring patients to a budesonide/formoterol fumarate fixed- dose combination therapy.

The benefits of inhaled budesonide therapy would normally minimise the need for oral steroids, but patients transferring from oral steroids may remain at risk of impaired adrenal reserve for a considerable time. Recovery may take a considerable amount of time after cessation of oral steroid therapy and hence oral steroid-dependent patients transferred to inhaled budesonide may remain at risk from impaired adrenal function for some considerable time. In such circumstances hypothalamic pituitary adrenocortical (HPA) axis function should be monitored regularly.

### High dose corticosteroids

The prolonged treatment with high doses of inhaled corticosteroids, particularly higher than recommended doses, may also result in clinically significant adrenal suppression. Therefore additional systemic corticosteroid cover should be considered during periods of stress such as severe infections or elective surgery. Rapid reduction in the dose of steroids can induce acute adrenal crisis. Symptoms and signs which might be seen in acute adrenal crisis may be somewhat vague but may include anorexia, abdominal pain, weight loss, tiredness, headache, nausea, vomiting, decreased level of consciousness, seizures, hypotension and hypoglycaemia.

Treatment with supplementary systematic steroids or inhaled budesonide should not be stopped abruptly.

### Transfer from oral therapy

During transfer from oral therapy to a budesonide/formoterol fumarate fixed-dose combination therapy, a generally lower systemic steroid action will be experienced which may result in the appearance of allergic or arthritic symptoms such as rhinitis, eczema and muscle and joint pain. Specific treatment should be initiated for these conditions. A general insufficient glucocorticosteroid effect should be suspected if, in rare cases, symptoms such as tiredness, headache, nausea and vomiting should occur. In these cases a temporary increase in the dose of oral glucocorticosteroids is sometimes necessary.

### Oral infections

To minimise the risk of oropharyngeal candida infection, the patient should be instructed to rinse their mouth out with water after inhaling the dose. If oropharyngeal thrush occurs, patients should also rinse their mouth with water after the as-needed inhalations.

### Pneumonia in patients with COPD

An increase in the incidence of pneumonia, including pneumonia requiring hospitalisation, has been observed in patients with COPD receiving inhaled corticosteroids. There is some evidence of an increased risk of pneumonia with increasing steroid dose but this has not been demonstrated conclusively across all studies.

There is no conclusive clinical evidence for intra-class differences in the magnitude of the pneumonia risk among inhaled corticosteroid products.

Physicians should remain vigilant for the possible development of pneumonia in patients with COPD as the clinical features of such infections overlap with the symptoms of COPD exacerbations.

Risk factors for pneumonia in patients with COPD include current smoking, older age, low body mass index (BMI) and severe COPD.

#### Interaction with other medicinal products

Concomitant treatment with itraconazole, ritonavir or other potent CYP3A4 inhibitors should be avoided (see section 4.5). If this is not possible the time interval between administrations of the interacting medicinal products should be as long as possible. In patients using potent CYP3A4 inhibitors, a budesonide/formoterol fumarate fixed-dose combination is not recommended.

#### Caution with special diseases

A fixed-dose combination of budesonide and formoterol fumarate dihydrate should be administered with caution in patients with thyrotoxicosis, phaeochromocytoma, diabetes mellitus, untreated hypokalaemia, hypertrophic obstructive cardiomyopathy, idiopathic subvalvular aortic stenosis, severe hypertension, aneurysm or other severe cardiovascular disorders, such as ischaemic heart disease, tachyarrhythmias or severe heart failure.

Caution should be observed when treating patients with prolongation of the QTc-interval. Formoterol itself may induce prolongation of the QTc-interval.

The need for, and dose of inhaled corticosteroids should be re-evaluated in patients with active or quiescent pulmonary tuberculosis, fungal and viral infections in the airways.

Additional blood glucose controls should be considered in diabetic patients.

#### $\beta_2$ adrenoreceptor agonists

Potentially serious hypokalaemia may result from high doses of  $\beta_2$  adrenoreceptor agonists. Concomitant treatment of  $\beta_2$  adrenoreceptor agonists with medicinal products which can induce hypokalaemia or potentiate a hypokalaemic effect, e.g. xanthine-derivatives, steroids and diuretics, may add to a possible hypokalaemic effect of the  $\beta_2$  adrenoreceptor agonist.

Treatment with  $\beta_2$  adrenoreceptor agonists may result in an increase in blood levels of insulin, free fatty acids, glycerol and ketone bodies.

Particular caution is recommended in unstable asthma with variable use of rescue bronchodilators, in acute severe asthma as the associated risk may be augmented by hypoxia and in other conditions when the likelihood for hypokalaemia is increased. It is recommended that serum potassium levels are monitored during these circumstances.

## Excipients

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal products.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Pharmacokinetic interactions

Potent inhibitors of CYP3A4 (e.g. ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin, telithromycin, nefazodone and HIV protease inhibitors) are likely to markedly increase plasma levels of budesonide and concomitant use should be avoided. If this is not possible the time interval between administration of the inhibitor and budesonide should be as long as possible (see section 4.4).

The potent CYP3A4 inhibitor ketoconazole, 200 mg once daily, increased plasma levels of concomitantly orally administered budesonide (single dose 3 mg) on average six-fold. When ketoconazole was administered 12 hours after budesonide the concentration was on average increased only three-fold showing that separation of the administration times can reduce the increase in plasma levels. Limited data about this interaction for high-dose inhaled budesonide indicates that marked increases in plasma levels (on average four fold) may occur if itraconazole, 200 mg once daily, is administered concomitantly with inhaled budesonide (single dose of 1000 micrograms).

Co-treatment with CYP3A inhibitors, including cobicistat-containing products, is expected to increase the risk of systemic side-effects. The combination should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects.<sup>1</sup>

### Pharmacodynamic interactions

$\beta$  adrenergic blockers can weaken or inhibit the effect of formoterol. A fixed-dose combination therapy of budesonide and formoterol fumarate dihydrate should therefore not be given together with  $\beta$  adrenergic blockers (including eye drops) unless there are compelling reasons.

Concomitant treatment with quinidine, disopyramide, procainamide, phenothiazines, antihistamines (terfenadine) and tricyclic antidepressants can prolong the QTc-interval and increase the risk of ventricular arrhythmias.

In addition L-Dopa, L-thyroxine, oxytocin and alcohol can impair cardiac tolerance towards  $\beta_2$  sympathomimetics.

Concomitant treatment with monoamine oxidase inhibitors including medicinal products with similar properties such as furazolidone and procarbazine may precipitate hypertensive reactions.

There is an elevated risk of arrhythmias in patients receiving concomitant anaesthesia with halogenated hydrocarbons.

Concomitant use of other  $\beta$  adrenergic medicinal products and anticholinergic medicinal products can have a potentially additive bronchodilating effect.

Hypokalaemia may increase the disposition towards arrhythmias in patients who are treated with digitalis glycosides.

Budesonide and formoterol have not been observed to interact with any other medicinal products used in the treatment of asthma.

#### Paediatric population

Interaction studies have only been performed in adults.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

For a fixed-dose combination therapy of budesonide and formoterol fumarate dihydrate or the concomitant treatment with formoterol and budesonide, no clinical data on exposed pregnancies are available. Data from an embryo-fetal development study in the rat, showed no evidence of any additional effect from the combination.

There are no adequate data from use of formoterol in pregnant women. In animal studies formoterol has caused adverse reactions in reproduction studies at very high systemic exposure levels (see section 5.3).

Data on approximately 2000 exposed pregnancies indicate no increased teratogenic risk associated with the use of inhaled budesonide. In animal studies glucocorticosteroids have been shown to induce malformations (see section 5.3). This is not likely to be relevant for humans given recommended doses.

Animal studies have also identified an involvement of excess prenatal glucocorticoids in increased risks for intrauterine growth retardation, adult cardiovascular disease and permanent changes in glucocorticoid receptor density, neurotransmitter turnover and behaviour at exposures below the teratogenic dose range.

During pregnancy, a fixed-dose combination therapy of budesonide and formoterol fumarate dihydrate should only be used when the benefits outweigh the potential risks. The lowest effective dose of budesonide needed to maintain adequate asthma control should be used.

#### Breast-feeding

Budesonide is excreted in breast milk. However, at therapeutic doses no effects on the suckling child are anticipated. It is not known whether formoterol passes into human breast milk. In rats, small amounts of formoterol have been detected in maternal milk. Administration of a fixed-dose combination therapy of budesonide and formoterol fumarate dihydrate to women who are breast-feeding should only be considered if the expected benefit to the mother is greater than any possible risk to the child.

#### Fertility

There is no data available on the potential effect of budesonide on fertility. Animal reproduction studies with formoterol have shown a somewhat reduced fertility in male rats at high systemic exposure (see section 5.3).

## 4.7 Effects on ability to drive and use machines

GoResp Digihaler has no or negligible influence on the ability to drive and use machines.

## 4.8 Undesirable effects

### Summary of the safety profile

Since GoResp Digihaler contains both budesonide and formoterol, the same pattern of adverse reactions as reported for these substances may occur. No increased incidence of adverse reactions has been seen following concurrent administration of the two compounds. The most common adverse reactions are pharmacologically predictable adverse reactions of  $\beta_2$  adrenoceptor agonist therapy, such as tremor and palpitations. These tend to be mild and usually disappear within a few days of treatment. In a 3-year clinical trial with budesonide in COPD, skin bruises and pneumonia occurred at a frequency of 10% and 6%, respectively, compared with 4% and 3% in the placebo group ( $p < 0.001$  and  $p < 0.01$ , respectively).

### Tabulated list of adverse reactions

Adverse reactions, which have been associated with budesonide or formoterol, are given below and listed by system organ class and frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$ ,  $< 1/10$ ), uncommon ( $\geq 1/1,000$ ,  $< 1/100$ ), rare ( $\geq 1/10,000$ ,  $< 1/1,000$ ), very rare ( $< 1/10,000$ ) and not known (cannot be estimated from the available data).

| System Organ Class                 | Frequency | Adverse reaction  |
|------------------------------------|-----------|---|
| Infections and infestations        | Common    | Candida infections in the oropharynx, pneumonia (in COPD patients)  |
| Immune system disorders            | Rare      | Immediate and delayed hypersensitivity reactions, e.g. exanthema, urticaria, pruritus, dermatitis, angioedema and anaphylactic reaction |
| Endocrine disorders                | Very rare | Cushing's syndrome, adrenal suppression, growth retardation, decrease in bone mineral density   |
| Metabolism and nutrition disorders | Rare      | Hypokalaemia  |
|                                    | Very rare | Hyperglycaemia  |
| Psychiatric disorders              | Uncommon  | Aggression, psychomotor hyperactivity, anxiety, sleep disorders   |
|                                    | Very rare | Depression, behavioural changes (predominantly in children)   |
| Nervous system disorders           | Common    | Headache, tremor  |
|                                    | Uncommon  | Dizziness   |
|                                    | Very rare | Taste disturbances  |
| Eye disorders                      | Very rare | Cataract and glaucoma   |
|                                    | Uncommon  | Vision, blurred (see also section 4.4)  |
| Cardiac disorders                  | Common    | Palpitations  |
|                                    | Uncommon  | Tachycardia   |
|                                    | Rare      | Cardiac arrhythmias, e.g. atrial fibrillation, supraventricular tachycardia, extrasystoles  |
|                                    | Very rare | Angina pectoris. Prolongation of QTc-interval   |
| Vascular disorders                 | Very rare | Variations in blood pressure  |

|   |           |   |
|---|-----------|---|
| Respiratory, thoracic and mediastinal disorders | Common    | Mild irritation in the throat, coughing, Dysphonia including hoarseness |
|   | Rare      | Bronchospasm  |
|   | Very rare | Paradoxical bronchospasm  |
| Gastrointestinal disorders                      | Uncommon  | Nausea  |
| Skin and subcutaneous tissue disorders          | Uncommon  | Bruises   |
| Musculoskeletal and connective tissue disorders | Uncommon  | Muscle cramps   |

#### Description of selected adverse reactions

Candida infection in the oropharynx is due to active substance deposition. Advising the patient to rinse the mouth out with water after each dose will minimise the risk. Oropharyngeal Candida infection usually responds to topical anti-fungal treatment without the need to discontinue the inhaled corticosteroid.

Paradoxical bronchospasm may occur very rarely, affecting less than 1 in 10,000 people, with an immediate increase in wheezing and shortness of breath after dosing. Paradoxical bronchospasm responds to a rapid-acting inhaled bronchodilator and should be treated straightaway. GoResp Digihaler should be discontinued immediately, the patient should be assessed and an alternative therapy is instituted if necessary (see section 4.4).

Systemic effects of inhaled corticosteroids may occur, particularly at high doses prescribed for long periods. These effects are much less likely to occur than with oral corticosteroids. Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma. Increased susceptibility to infections and impairment of the ability to adapt to stress may also occur. Effects are probably dependent on dose, exposure time, concomitant and previous steroid exposure and individual sensitivity.

Treatment with  $\beta_2$  adrenoceptor agonists may result in an increase in blood levels of insulin, free fatty acids, glycerol and ketone bodies.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

An overdose of formoterol would likely lead to effects that are typical for  $\beta_2$  adrenoceptor agonists: tremor, headache, palpitations. Symptoms reported from isolated cases are tachycardia, hyperglycaemia, hypokalaemia, prolonged QTc-interval, arrhythmia, nausea and vomiting. Supportive and symptomatic treatment may be indicated. A dose of 90 micrograms administered during three hours in patients with acute bronchial obstruction raised no safety concerns.

Acute overdose with budesonide, even in excessive doses, is not expected to be a clinical problem. When used chronically in excessive doses, systemic glucocorticosteroid effects, such as hypercorticism and adrenal suppression, may appear.

If GoResp Digihaler therapy has to be withdrawn due to overdose of the formoterol component of the medicinal product, provision of appropriate inhaled corticosteroid therapy must be considered.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for obstructive airway diseases, adrenergics and other drugs for obstructive airway diseases.

ATC code: R03AK07

#### Mechanism of action and pharmacodynamic effects

GoResp Digihaler contains formoterol and budesonide, which have different modes of action and show additive effects in terms of reduction of asthma exacerbations. The mechanisms of action of the two substances respectively are discussed below.

##### *Budesonide*

Budesonide is a glucocorticosteroid which when inhaled has a dose-dependent anti-inflammatory action in the airways, resulting in reduced symptoms and fewer asthma exacerbations. Inhaled budesonide has less severe adverse reactions than systemic corticosteroids. The exact mechanism responsible for the anti-inflammatory effect of glucocorticosteroids is unknown.

##### *Formoterol*

Formoterol is a selective  $\beta_2$  adrenoceptor agonist that when inhaled results in rapid and long-acting relaxation of bronchial smooth muscle in patients with reversible airways obstruction. The bronchodilating effect is dose-dependent, with an onset of effect within 1-3 minutes. The duration of effect is at least 12 hours after a single dose.

#### Clinical efficacy and safety

##### *Asthma*

##### *Budesonide/formoterol maintenance therapy*

Clinical studies in adults have shown that the addition of formoterol to budesonide improved asthma symptoms and lung function, and reduced exacerbations.

In two 12-week studies the effect on lung function of budesonide/formoterol was equal to that of the free combination of budesonide and formoterol, and exceeded that of budesonide alone. All treatment arms used a short-acting  $\beta_2$  adrenoceptor agonist as needed. There was no sign of attenuation of the anti-asthmatic effect over time.

Two 12-week paediatric studies have been performed in which 265 children aged 6-11 years were treated with a maintenance dose of budesonide/formoterol (2 inhalations of 80 micrograms/4.5 micrograms/inhalation twice daily), and a short acting  $\beta_2$  adrenoceptor agonist as needed. In both studies, lung function was improved and the treatment was well tolerated compared to the corresponding dose of budesonide alone (see paediatric population sub-heading in section 4.2).

### COPD

In two 12-month studies, the effect on lung function and the rate of exacerbation (defined as courses of oral steroids and/or course of antibiotics and/or hospitalisations) in patients with moderate to severe COPD was evaluated. The inclusion criteria for both studies was prebronchodilator FEV<sub>1</sub> <50% predicted normal. Median post-bronchodilator FEV<sub>1</sub> at inclusion in the trials was 42% predicted normal.

The mean number of exacerbations per year (as defined above) was significantly reduced with budesonide/formoterol as compared with treatment with formoterol alone or placebo (mean rate 1.4 compared with 1.8-1.9 in the placebo/formoterol group). The mean number of days on oral corticosteroids/patient during the 12 months was slightly reduced in the budesonide/formoterol group (7-8 days/patient/year compared with 11-12 and 9-12 days in the placebo and formoterol groups, respectively). For changes in lung-function parameters, such as FEV<sub>1</sub>, budesonide/formoterol was not superior to treatment with formoterol alone.

### Peak Inspiratory Flow Rate through the Digihaler Device

The integrated eModule in Digihaler (a digital version of Spiromax) does not impact the metering or dosing functionality of the inhaler and its performance has been proven to be equivalent to that of Spiromax. Therefore the below findings from the study on Spiromax are also applicable to Digihaler.

A randomised, open-label placebo study was performed in children and adolescents with asthma (aged 6-17 years), adults with asthma (aged 18-45 years), adults with chronic obstructive pulmonary disease (COPD – aged >50 years) and healthy volunteers (aged 18-45 years) to evaluate the peak inspiratory flow rate (PIFR) and other related inhalation parameters following inhalation from a Spiromax device (containing placebo) compared with inhalation from an already marketed multi-dose dry powder inhaler device (containing placebo). The impact of enhanced training in dry powder inhaler inhalation technique on inhalation speed and volume was also assessed in these subject groups. The data from the study indicated that regardless of age and underlying disease severity, children, adolescents and adults with asthma as well as patients with COPD were able to achieve inspiratory flow rates through the Spiromax device that were similar to those generated through the marketed multi-dose dry powder inhaler device (see paediatric population sub-heading in section 4.2). The mean PIFR achieved by patients with asthma or COPD was over 60L/min, a flow rate at which both devices studied are known to deliver comparable amounts of drug to the lungs. Very few patients had PIFRs below 40L/min; when PIFRs were less than 40L/min there appeared to be no clustering by age or disease severity.

An open-label, single-visit study was conducted to demonstrate the accuracy of the Digihaler in measuring inhalation parameters in paediatric patients ages 4 to 17 years with asthma, adults 18 to 55 years of age with asthma (FEV<sub>1</sub><80% of predicted), and adults 55 years of age and older with COPD (FEV<sub>1</sub> <50% of predicted). Empty inhalers (no drug or excipient) were used by patients after receiving training in correct technique and the accuracy of the Digihaler was assessed by comparing the inhalation parameters, including PIF, recorded simultaneously

by the eModule and by an inhalation profile recorder (IPR) attached to the inhaler. Both the PIF and the inhalation volume results demonstrated the accuracy of the empty Digihaler. Except for two subjects with COPD, all other patients in this study could generate sufficient inspiratory flow rates (above 30 L/min) through the Digihaler for adequate dose delivery.

## **5.2 Pharmacokinetic properties**

### **Absorption**

The fixed-dose combination of budesonide and formoterol, and the corresponding monoproducts have been shown to be bioequivalent with regard to systemic exposure of budesonide and formoterol, respectively. In spite of this, a small increase in cortisol suppression was seen after administration of fixed-dose combination compared to the monoproducts. The difference is considered not to have an impact on clinical safety.

There was no evidence of pharmacokinetic interactions between budesonide and formoterol.

Pharmacokinetic parameters for the respective substances were comparable after the administration of budesonide and formoterol as monoproducts or as the fixed-dose combination. For budesonide, AUC was slightly higher, rate of absorption more rapid and maximal plasma concentration higher after administration of the fixed combination. For formoterol, maximal plasma concentration was similar after administration of the fixed combination. Inhaled budesonide is rapidly absorbed and the maximum plasma concentration is reached within 30 minutes after inhalation. In studies, mean lung deposition of budesonide after inhalation via the powder inhaler ranged from 32% to 44% of the delivered dose. The systemic bioavailability is approximately 49% of the delivered dose. In children 6-16 years of age the lung deposition falls in the same range as in adults for the same given dose (see paediatric population sub-heading in section 4.2). The resulting plasma concentrations were not determined.

Inhaled formoterol is rapidly absorbed and the maximum plasma concentration is reached within 10 minutes after inhalation. In studies the mean lung deposition of formoterol after inhalation via the powder inhaler ranged from 28% to 49% of the delivered dose. The systemic bioavailability is about 61% of the delivered dose.

### **Distribution**

Plasma protein binding is approximately 50% for formoterol and 90% for budesonide. Volume of distribution is about 4 L/kg for formoterol and 3 L/kg for budesonide. Formoterol is inactivated via conjugation reactions (active O-demethylated and deformedylated metabolites are formed, but they are seen mainly as inactivated conjugates). Budesonide undergoes an extensive degree (approximately 90%) of biotransformation on first passage through the liver to metabolites of low glucocorticosteroid activity. The glucocorticosteroid activity of the major metabolites, 6-beta-hydroxy-budesonide and 16-alfa-hydroxy-prednisolone, is less than 1% of that of budesonide. There are no indications of any metabolic interactions or any displacement reactions between formoterol and budesonide.

### **Elimination**

The major part of a dose of formoterol is transformed by liver metabolism followed by renal elimination. After inhalation, 8% to 13% of the delivered dose of formoterol is excreted unmetabolised in the urine. Formoterol has a high systemic clearance (approximately 1.4 L/min) and the terminal elimination half-life averages 17 hours.

Budesonide is eliminated via metabolism mainly catalysed by the enzyme CYP3A4. The metabolites of budesonide are eliminated in urine as such or in conjugated form. Only negligible amounts of unchanged budesonide have been detected in the urine. Budesonide has a high systemic clearance (approximately 1.2 L/min) and the plasma elimination half-life after i.v. dosing averages 4 hours.

#### Pharmacokinetic/pharmacodynamic relationship(s)

The pharmacokinetics of budesonide or formoterol in children and patients with renal failure are unknown. The exposure of budesonide and formoterol may be increased in patients with liver disease.

#### Budesonide/Formoterol Spiromax pharmacokinetic profile

In pharmacokinetic studies with and without a charcoal blockage, Budesonide/Formoterol Spiromax was evaluated by comparing it with an alternative authorised fixed-dose combination inhaled product containing the same active substances, budesonide and formoterol and has been shown to be equivalent in both systemic exposure (safety) and pulmonary deposition (efficacy).

#### Linearity/non-linearity

Systemic exposure for both budesonide and formoterol correlates in a linear fashion to administered dose.

The results of these studies are also applicable for GoResp Digihaler, as it contains the same active substances, budesonide and formoterol fumarate dihydrate, and the performance of Digihaler has been proven to be equivalent to that of Spiromax.

### **5.3 Preclinical safety data**

The toxicity observed in animal studies with budesonide and formoterol, given in combination or separately, were effects associated with exaggerated pharmacological activity.

In animal reproduction studies, corticosteroids such as budesonide have been shown to induce malformations (cleft palate, skeletal malformations). However, these animal experimental results do not seem to be relevant in humans at the recommended doses. Animal reproduction studies with formoterol have shown a somewhat reduced fertility in male rats at high systemic exposure and implantation losses as well as decreased early postnatal survival and birth weight at considerably higher systemic exposures than those reached during clinical use. However, these animal experimental results do not seem to be relevant in humans.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate.

### **6.2 Incompatibilities**

Not applicable.

### 6.3 Shelf life

3 years.

After opening the foil wrap: 6 months.

### 6.4 Special precautions for storage

Do not store above 25°C.

Keep the mouthpiece cover closed after removal of the foil wrap.

### 6.5 Nature and contents of container

The inhaler is white (with an integrated electronic Module, eModule, at the top) and a semi-transparent wine red mouthpiece cover. The top of the inhaler there is a QR code that can be scanned by a mobile device using the Digihaler mobile application (once downloaded) in order to connect the inhaler to the App (via Bluetooth®). The drug/mucosal contact parts of the inhaler are made of acrylonitrile butadiene styrene (ABS), polyethylene (PE), and polypropylene (PP). Each inhaler contains 90 doses and is foil-wrapped.

Pack sizes of 1, 2 or 3 inhalers.

Not all pack-sizes may be marketed.

The below symbols appear on the inhaler (laser printed) or on the labelling:



Read the IFU before use symbol



Degree of protection of applied parts against electric shock

FCC ID: 2AJVSPDR0000401 Federal Communications Commission Identification



Battery disposal



Bluetooth transmitter symbol

### 6.6 Special precautions for disposal and other handling

GoResp Digihaler contains a lithium-manganese dioxide battery and must not be thrown away via household waste. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Do not immerse GoResp Digihaler in water.

## **7      MARKETING AUTHORISATION HOLDER**

Teva UK Limited,  
Ridings Point,  
Whistler Drive,  
Castleford, WF10 5HX,  
United Kingdom

## **8      MARKETING AUTHORISATION NUMBER(S)**

PLGB 00289/2502

## **9      DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

01/01/2021

## **10     DATE OF REVISION OF THE TEXT**

01/03/2024

The Bluetooth® word mark and logos are registered trademarks owned by the Bluetooth SIG, Inc. and any use of such marks by Teva Respiratory, LLC is under license.


This device has been tested to EN IEC 60601-1-2:2015 +A1:2021 to the requirements for home health care for both emissions and immunity. *Note that portable RF (Radio Frequency) communications equipment, including peripherals such as antenna cables and external antenna, should be used no closer than 30cm (12 inches) to the inhaler whilst the inhaler is being used. Otherwise, registering the use of the inhaler might be delayed or not recorded. You may need to open the app later to see such notifications.*

The inhaler and its associated app running on a smart device are intended for use in the home healthcare environment. *Note that the inhaler can be used in an aircraft, but the smart device should be in “aircraft/flight mode” when required by the cabin crew.*

Patients should not bring the inhaler(s) into an MRI or CT/X-ray scan room. The inhaler should not be exposed to other known sources of electromagnetic interference (such as diathermy). The inhaler can be taken through standard security systems, such as metal detectors.

### **IP22**

Protected against solid objects over 12.5 mm and direct sprays of water up to 15° from the vertical



**UK  
CA**

For England, Scotland and Wales Hereby, TEVA declares that the Bluetooth radio equipment contained in this product complies with the requirements of the UK Radio Equipment Regulations (2017). The full text of the UKCA declaration of conformity is available at the following internet address: [www.tevadigihaler.co.uk](http://www.tevadigihaler.co.uk)