



Tacrolimus 0.1% Ointment

tacrolimus monohydrate

Package Leaflet: Information for the user

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Tacrolimus 0.1% Ointment is and what it is used for
2. What you need to know before you use Tacrolimus 0.1% Ointment
3. How to use Tacrolimus 0.1% Ointment
4. Possible side effects
5. How to store Tacrolimus 0.1% Ointment
6. Contents of the pack and other information

1 What Tacrolimus 0.1% Ointment is and what it is used for

The active substance of Tacrolimus 0.1 % Ointment, tacrolimus monohydrate, is an immunomodulating agent.

Tacrolimus 0.1% Ointment is used to treat moderate to severe atopic dermatitis (eczema) in adults who are not adequately responsive to or are intolerant of conventional therapies such as topical corticosteroids.

Once moderate to severe atopic dermatitis is cleared or almost cleared after up to 6 weeks treatment of a flare, and if you are experiencing frequent flares (i.e. 4 or more per year), it may be possible to prevent flares coming back or prolong the time you are free from flares by using Tacrolimus 0.1% Ointment twice weekly.

In atopic dermatitis, an over-reaction of the skin's immune system causes skin inflammation (itchiness, redness, dryness). Tacrolimus 0.1 % Ointment alters the abnormal immune response and relieves the skin inflammation and the itch.

2 What you need to know before you use Tacrolimus 0.1% Ointment

Do not use Tacrolimus 0.1 % Ointment

- If you are allergic to tacrolimus or any of the other ingredients of this medicine (listed in section 6) or to macrolide antibiotics (e.g. azithromycin, clarithromycin, erythromycin).

Warnings and precautions

Talk to your doctor if you:

- have **liver failure**
- have any **skin malignancies** (tumours) or if you have a **weakened immune system** (immuno-compromised) whatever the cause
- have an **inherited skin barrier disease** such as Netherton's syndrome, lamellar ichthyosis (extensive scaling of the skin due to a thickening of the outer layer of the skin), or if you have an inflammatory skin disease such as pyoderma gangrenosum, or if you suffer from **generalised erythroderma** (inflammatory reddening and scaling of the entire skin)
- have a cutaneous Graft Versus Host Disease (an immune reaction of the skin which is a common

complication in patients who have undergone a bone marrow transplant)

- have **swollen lymph nodes** at initiation of treatment. If your lymph nodes become swollen during treatment with Tacrolimus 0.1% Ointment, consult your doctor
- have **infected lesions**. Do not apply the ointment to infected lesions.
- notice any **change to the appearance of your skin**, please inform your physician.
- the safety of using tacrolimus ointment for a long time is not known. A very small number of people who have used tacrolimus ointment have had malignancies (for example, skin or lymphoma). However, a link to Tacrolimus 0.1% Ointment treatment has not been shown.
- avoid exposing the skin to long periods of sunlight or artificial sunlight such as tanning beds. If you spend time outdoors after applying Tacrolimus 0.1% Ointment, use a sunscreen and wear loose fitting clothing that protects the skin from the sun. In addition, ask your doctor for advice on other appropriate sun protection methods. If you are prescribed light therapy, inform your doctor that you are using Tacrolimus 0.1% Ointment as it is not recommended to use Tacrolimus 0.1% Ointment and light therapy at the same time.
- if your doctor tells you to use Tacrolimus 0.1% Ointment twice weekly to keep your atopic dermatitis cleared, your condition should be reviewed by your doctor at least every 12 months, even if it remains under control. In children, maintenance treatment should be suspended after 12 months, to assess whether the need for continued treatment still exists.
- do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.
- it is recommended to use Tacrolimus Ointment at the lowest possible strength, at the lowest frequency and for the shortest possible duration necessary. This decision should be based on your doctor's assessment of how your eczema responds to Tacrolimus Ointment.

Children

- Tacrolimus 0.1 % Ointment is **not approved for children younger than 16 years of age**. Therefore it should not be used in this age group. Please consult your doctor.
- The effect of treatment with Tacrolimus 0.1% Ointment on the developing immune system in children, especially the young, has not been established.

Other medicines, cosmetics and Tacrolimus 0.1% Ointment

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

You may use moisturising creams and lotions during treatment with Tacrolimus 0.1% Ointment but these products should not be used within two hours of applying Tacrolimus 0.1% Ointment.

The use of tacrolimus at the same time as other preparations to be used on the skin or while taking oral corticosteroids (e.g. cortisone) or medicines which affect the immune system has not been studied.

Tacrolimus 0.1% Ointment with alcohol

While using Tacrolimus 0.1% Ointment, drinking alcohol may cause the skin or face to become flushed or red and feel hot.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

3 How to use Tacrolimus 0.1% Ointment

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- apply Tacrolimus 0.1% Ointment as a thin layer to affected areas of your skin
- tacrolimus 0.1% Ointment may be used on most parts of the body, including the face and neck and in the creases of your elbows and knees.
- avoid using the ointment inside your nose or mouth or in your eyes. If the ointment gets on any of these areas, it should be thoroughly wiped off and/or rinsed off with water.
- do not cover the skin being treated with bandages, wraps or other wound dressing
- wash your hands after applying Tacrolimus 0.1% Ointment unless your hands are also being treated
- before applying Tacrolimus 0.1% Ointment after a bath or shower, be sure your skin is completely dry.

Adults (16 years of age and older)

Two strengths of tacrolimus ointment (tacrolimus 0.03% ointment and tacrolimus 0.1% ointment) are available for adult patients (16 years of age and older). Your doctor will decide which strength is best for you.

Usually, treatment is started with Tacrolimus 0.1% Ointment twice a day, once in the morning and once in the evening, until the eczema has cleared. Depending on the response of your eczema your doctor will decide if the frequency of application can be reduced or the lower strength, tacrolimus 0.03% ointment, can be used.

Treat each affected region of your skin until the eczema has gone away. Improvement is usually seen within one week. If you do not see any improvement after two weeks, see your doctor about other possible treatments.

You may be told by your doctor to use Tacrolimus 0.1% Ointment twice weekly once your atopic dermatitis has cleared or almost cleared. Tacrolimus 0.1 % Ointment should be applied once a day twice weekly (e.g. Monday and Thursday) to areas of your body commonly affected by atopic dermatitis. There should be 2–3 days without Tacrolimus 0.1% Ointment treatment between applications.

If symptoms reappear you should use Tacrolimus 0.1 % Ointment twice daily as outlined above and arrange to see your doctor to review your treatment.

If you accidentally swallow some Tacrolimus 0.1% Ointment

If you accidentally swallow the ointment, consult your doctor or pharmacist as soon as possible. Do not try to induce vomiting.

If you forget to use Tacrolimus 0.1% Ointment

If you forget to apply the ointment at the scheduled time, do it as soon as you remember and then continue as before.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common: may affect more than 1 in 10 people

- burning sensation and itching
- These symptoms are usually mild to moderate and generally go away within one week of using Tacrolimus 0.1% Ointment.

Common: may affect up to 1 in 10 people

- redness
- feeling of warmth
- pain

- increased skin sensitivity (especially to hot and cold)
- skin tingling and irritation
- rash
- local skin infection regardless of specific cause including but not limited to: inflamed or infected hair follicles, cold sores, generalised herpes simplex infections
- facial flushing or skin irritation after drinking alcohol is also common

Uncommon: may affect fewer than 1 in 100 people

- acne

Following twice-weekly treatment application site infections have been reported in adults.

Rosacea (facial redness), rosacea-like dermatitis, lentigo (presence of flat brown spots on the skin), oedema at the application site and herpes eye infections have been reported during post-marketing experience.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Tacrolimus 0.1% Ointment

Keep this medicine out of the sight and reach of children.

Do not use Tacrolimus 0.1% Ointment after the expiry date which is stated on the tube and carton after EXP. The expiry date refers to the last day of that month. Do not store above 25°C.

Discard open tubes 90 days after opening, even if they are not empty. They should not be kept for future use.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6 Contents of the pack and other information

What Tacrolimus 0.1% Ointment contains

- The active substance is tacrolimus monohydrate. One gram of Tacrolimus 0.1% Ointment contains 1.0 mg tacrolimus (as tacrolimus monohydrate).
- The other ingredients are paraffin, white soft, paraffin liquid, propylene carbonate, beeswax, white, paraffin hard.

What Tacrolimus 0.1% Ointment looks like and contents of the pack

Tacrolimus 0.1% Ointment is a white to slightly yellowish ointment. It is supplied in tubes containing 10, 30 or 60 grams of ointment. Not all pack sizes may be marketed.

Marketing Authorisation Holder

Teva UK Limited, Ridings Point, Whistler Drive, Castleford, WF10 5HX, United Kingdom

Manufacturer

LABORATORI FUNDACIÓ DAU, Pol. Ind. Consorci Zona Franca, c/ C, 12-14 Barcelona, 08040, Spain

This leaflet was last revised in November 2024.

PL 00289/2178

teva

215276_s1
INP010
10.00000.0.000000