

## Package leaflet: Information for the patient

# Dimethyl Fumarate 120 mg and 240 mg Gastro-resistant Hard Capsules

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Dimethyl Fumarate Gastro-resistant Hard Capsules are and what they are used for
2. What you need to know before you take Dimethyl Fumarate Gastro-resistant Hard Capsules
3. How to take Dimethyl Fumarate Gastro-resistant Hard Capsules
4. Possible side effects
5. How to store Dimethyl Fumarate Gastro-resistant Hard Capsules
6. Contents of the pack and other information

## 1. What Dimethyl Fumarate Gastro-resistant Hard Capsules are and what they are used for

### What Dimethyl Fumarate is

Dimethyl Fumarate Gastro-resistant Hard Capsules are a medicine that contains the active substance **dimethyl fumarate**.

### What Dimethyl Fumarate is used for

**Dimethyl Fumarate is used to treat relapsing-remitting multiple sclerosis (MS) in patients aged 13 years and older.**

MS is a long-term condition that affects the central nervous system (CNS), including the brain and the spinal cord. Relapsing-remitting MS is characterised by repeated attacks (relapses) of nervous system symptoms. Symptoms vary from patient to patient, but typically include walking difficulties, feeling off balance and visual difficulties (e.g. blurred or double vision). These symptoms may disappear completely when the relapse is over, but some problems may remain.

### How Dimethyl Fumarate works

Dimethyl Fumarate seems to work by stopping the body's defence system from damaging your brain and spinal cord. This may also help to delay future worsening of your MS.

## 2. What you need to know before you take Dimethyl Fumarate Gastro-resistant Hard Capsules

### Do not take Dimethyl Fumarate:

- **if you are allergic to dimethyl fumarate** or any of the other ingredients of this medicine (listed in section 6)
- **if you are suspected to suffer from a rare brain infection called progressive multifocal leukoencephalopathy (PML) or if PML has been confirmed.**

### Warnings and precautions

Dimethyl Fumarate may affect your **white blood cell counts**, your **kidneys** and **liver**. Before you start Dimethyl Fumarate, your doctor will do a blood test to count the number of your white blood cells and will check that your kidneys and liver are working properly. Your doctor will test these periodically during treatment. If your number of white blood cells decreases during treatment, your doctor may consider additional tests or discontinue your treatment.

**Talk to your doctor** before taking Dimethyl Fumarate if you have:

- severe **kidney** disease
- severe **liver** disease
- a disease of the **stomach** or **bowel**
- a serious **infection** (such as pneumonia).

Herpes zoster (shingles) may occur with Dimethyl Fumarate treatment. In some cases, serious complications have occurred. **You should inform your doctor** immediately if you suspect you have any symptoms of shingles.

If you believe your MS is getting worse (e.g. weakness or visual changes) or if you notice any new symptoms, talk to your doctor straight away because these may be the symptoms of a rare brain infection called PML. PML is a serious condition that may lead to severe disability or death.

A rare but serious kidney disorder called Fanconi Syndrome has been reported with a medicine containing dimethyl fumarate, in combination with other fumaric acid esters, used to treat psoriasis (a skin disease). If you notice you are

## Serious side effects

Dimethyl Fumarate may lower lymphocyte counts (a type of white blood cell). Having a low white blood cell count can increase your risk of infection, including the risk of a rare brain infection called progressive multifocal leukoencephalopathy (PML). PML may lead to severe disability or death. PML has occurred after 1 to 5 years of treatment and so your physician should continue to monitor your white blood cells throughout your treatment, and you should remain observant of any potential symptoms of PML as described below. The risk of PML may be higher if you have previously taken a medicine impairing the functionality of your body's immune system.

The symptoms of PML may be similar to an MS relapse. Symptoms may include new or worsening weakness on one side of the body; clumsiness; changes in vision, thinking, or memory; or confusion or personality changes, or speech and communication difficulties lasting for more than several days. Therefore, if you believe your MS is getting worse or if you notice any new symptoms while you are on Dimethyl Fumarate treatment, it is very important that you speak to your doctor as soon as possible. Also speak with your partner or caregivers and inform them about your treatment. Symptoms might arise that you might not become aware of by yourself.

**Call your doctor straight away if you experience any of these symptoms.**

## Severe allergic reactions

The frequency of severe allergic reactions cannot be estimated from the available data (not known).

Reddening of the face or body (flushing) is a very common side effect. However, should flushing be accompanied by a red rash or hives **and** you get any of these symptoms:

- swelling of the face, lips, mouth or tongue (angioedema)
- wheezing, difficulty breathing or shortness of breath (dyspnoea, hypoxia)
- dizziness or loss of consciousness (hypotension).

then this may represent a severe allergic reaction (anaphylaxis).

**Stop taking Dimethyl Fumarate and call a doctor straight away.**

## Other side effects

**Very common** (may affect more than 1 in 10 people)

- reddening of the face or body feeling warm, hot, burning or itchy (flushing)
- loose stools (diarrhoea)
- feeling sick (nausea)
- stomach pain or stomach cramps.

**Taking your medicine with food can help to reduce the side effects above.**

Substances called ketones, which are naturally produced in the body, very commonly show up in urine tests while taking Dimethyl Fumarate.

Talk to your doctor about how to manage these side effects. Your doctor may reduce your dose. Do not reduce your dose unless your doctor tells you to.

**Common** (may affect up to 1 in 10 people)

- inflammation of the lining of the intestines (gastroenteritis)
- being sick (vomiting)
- indigestion (dyspepsia)
- inflammation of the lining of the stomach (gastritis)
- gastrointestinal disorder
- burning sensation
- hot flush, feeling hot
- itchy skin (pruritus)
- rash
- pink or red blotches on the skin (erythema)
- hair loss (alopecia).

Side effects which may show up in your blood or urine tests

- low levels of white blood cells (lymphopenia, leucopenia) in the blood. Reduced white blood cells could mean your body is less able to fight an infection. If you have a serious infection (such as pneumonia), talk to your doctor immediately.
- proteins (albumin) in urine
- increase in levels of liver enzymes (ALT, AST) in the blood.

**Uncommon** (may affect up to 1 in 100 people)

- allergic reactions (hypersensitivity)
- reduction in blood platelets.

**Not known** (frequency cannot be estimated from the available data)

- liver inflammation and increase in levels of liver enzymes (ALT or AST in combination with bilirubin)
- herpes zoster (shingles) with symptoms such as blisters, burning, itching or pain of the skin, typically on one side of the upper body or the face, and other symptoms, like fever and weakness in the early stages of infection, followed by numbness, itching or red patches with severe pain
- runny nose (rhinorrhoea).

## Children (13 years of age and above) and adolescents

The side effects listed above also apply to children and adolescents.

Some side effects were reported more frequently in children and adolescents than in adults, e.g., headache, stomach pain or stomach cramps, being sick (vomiting), throat pain, cough and painful menstrual periods.

## Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Dimethyl Fumarate Gastro-resistant Hard Capsules

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister or bottle label and the carton after "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special temperature storage conditions.

Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Dimethyl Fumarate 120 mg and 240 mg Gastro-resistant Hard Capsules contains

The active substance is dimethyl fumarate.

### Dimethyl Fumarate 120 mg Gastro-resistant Hard Capsules

Each hard gastro-resistant capsule contains 120 mg of dimethyl fumarate.

### Dimethyl Fumarate 240 mg Gastro-resistant Hard Capsules

Each hard gastro-resistant capsule contains 240 mg of dimethyl fumarate.

The other ingredients are:

- Capsule content: cellulose, microcrystalline, crospovidone, talc, povidone, silica colloidal anhydrous, magnesium stearate, triethyl citrate, methacrylic acid/ethylacrylate copolymer (1:1), hypromellose, titanium dioxide (E 171), triacetin.
- Capsule shell: gelatin, titanium dioxide (E 171), brilliant blue FCF-FD&C Blue 1 (E 133), yellow iron oxide (E 172).
- Capsule printing ink: shellac, potassium hydroxide, propylene glycol (E 1520), black iron oxide (E 172), strong ammonia solution.

### What Dimethyl Fumarate 120 mg and 240 mg Gastro-resistant Hard Capsules looks like and contents of the pack

Dimethyl Fumarate 120 mg Gastro-resistant Hard Capsules (gastro-resistant capsules): green cap and white body, capsule shell of 21.4 mm, imprinted in black ink with "DMF 120" on the body containing white to off-white minitables.

Dimethyl Fumarate 240 mg Gastro-resistant Hard Capsules (gastro-resistant capsules): green cap and body, capsule shell of 23.2 mm, imprinted in black ink with "DMF 240" on the body containing white to off-white minitables.

OPA/Alu/PVC//Alu blisters or OPA/Alu/PVC//Alu unit dose blister.

### Dimethyl Fumarate 120 mg Gastro-resistant Hard Capsules

Pack sizes:

14 capsules (blisters).

14 x 1 capsules (unit dose perforated blisters).

### Dimethyl Fumarate 240 mg Gastro-resistant Hard Capsules

Pack sizes:

56 capsules (blisters).

56 x 1 capsules (unit dose perforated blisters).

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Teva UK Limited, Ridings Point, Whistler Drive, Castleford, WF10 5HX, United Kingdom

### Manufacturer

Pharmadox Healthcare Limited, Kw20a Kordin Industrial Estate, Paola, PLA 3000, Malta

**This leaflet was last revised in November 2024.**

PLGB 00289/2576

PLGB 00289/2577