

Candesartan cilexetil 8 mg Tablets
Candesartan cilexetil 16 mg Tablets
Candesartan cilexetil 32 mg Tablets

Package leaflet: Information for the user

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Candesartan cilexetil is and what it is used for
2. What you need to know before you take Candesartan cilexetil
3. How to take Candesartan cilexetil
4. Possible side effects
5. How to store Candesartan cilexetil
6. Contents of the pack and other information

1 What Candesartan cilexetil is and what it is used for

The name of your medicine is Candesartan cilexetil. The active ingredient is candesartan cilexetil. This belongs to a group of medicines called Angiotensin II receptor antagonists. It works by making your blood vessels relax and widen. This helps to lower your blood pressure. It also makes it easier for your heart to pump blood to all parts of your body.

This medicine is used for:

- treating high blood pressure (hypertension) in adult patients and in children and adolescents aged 6 to <18 years.
- treating adult heart failure patients with reduced heart muscle function when Angiotensin Converting Enzyme (ACE) inhibitors cannot be used or in addition to ACE-inhibitors when symptoms persist despite treatment and mineralocorticoid receptor antagonists (MRA) cannot be used. (ACE-inhibitors and MRAs are medicines used to treat heart failure).

2 What you need to know before you take Candesartan cilexetil

Do not take Candesartan cilexetil

- if you are allergic to Candesartan cilexetil or any of the other ingredients of this medicine (listed in section 6).
- if you are more than 3 months pregnant (it is also better to avoid this medicine in early pregnancy – see section 2: Pregnancy and breast-feeding).
- if you have severe liver disease or biliary obstruction (a problem with the drainage of the bile from the gall bladder).
- if the patient is a child under 1 year of age.
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren

If you are not sure if any of these apply to you, talk to your doctor or pharmacist before taking this medicine.

Warnings and precautions

Talk to your doctor or pharmacist before taking Candesartan cilexetil:

- if you have heart, liver or kidney problems, or are on dialysis
- if you have recently had a kidney transplant
- if you are vomiting, have recently had severe vomiting, or have diarrhoea
- if you have a disease of the adrenal gland called Conn's syndrome (also called primary hyperaldosteronism)
- if you have low blood pressure
- if you have ever had a stroke
- if you must tell your doctor if you think you are (or might become) pregnant. This medicine is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see section 2: Pregnancy and breast-feeding).
- if you are taking any of the following medicines used to treat high blood pressure:
 - an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.
 - aliskiren
- if you are taking an ACE-inhibitor together with a medicine which belongs to the class of medicines known as mineralocorticoid receptors antagonists (MRA). These medicines are for the treatment of heart failure (see 'Other medicines and Candesartan cilexetil').

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals. See also information under the heading 'Do not take Candesartan cilexetil'.

Your doctor may want to see you more often and do some tests if you have any of these conditions.

Talk to your doctor if you experience abdominal pain, nausea, vomiting or diarrhoea after taking Candesartan cilexetil.

Your doctor will decide on further treatment. Do not stop taking Candesartan cilexetil on your own.

If you are going to have an operation, tell your doctor or dentist that you are taking Candesartan cilexetil. This is because this medicine when combined with some anaesthetics, may cause an excessive drop in blood pressure.

Children and adolescents

Candesartan has been studied in children. For more information, talk to your doctor. Candesartan must not be given to children under 1 year of age due to the potential risk to the developing kidneys.

Other medicines and Candesartan Cilexetil

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. Candesartan cilexetil can affect the way some other medicines work and some medicines can have an effect on Candesartan cilexetil. If you are using certain medicines, your doctor may need to do blood tests from time to time.

In particular, tell your doctor if you are using any of the following medicines:

- other medicines to help lower your blood pressure, including beta-blockers, diazoxide and ACE inhibitors such as enalapril, captopril, lisinopril or ramipril.
- non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen, naproxen, diclofenac, celecoxib or etoricoxib (medicines to relieve pain and inflammation).
- acetylsalicylic acid (if you are taking more than 3 g each day) (medicine to relieve pain and inflammation).
- potassium supplements or salt substitutes containing potassium (medicines that increase the amount of potassium in your blood).
- heparin (a medicine for thinning the blood).
- co-trimoxazole (an antibiotic medicine) also known as trimethoprim/sulfamethoxazole.
- water tablets (diuretics).
- lithium (a medicine for mental health problems).

Your doctor may need to change your dose and/or to take other precautions:

- If you are taking an ACE-inhibitor or aliskiren (see also information under the headings 'Do not take Candesartan cilexetil' and 'Warnings and precautions')
- If you are being treated with an ACE-inhibitor together with certain other medicines to treat your heart failure, which are known as mineralocorticoid receptors antagonists (MRA) (for example spironolactone, eplerenone).

Candesartan cilexetil with food, drink and alcohol

- You can take this medicine with or without food.
- When you are prescribed Candesartan cilexetil, discuss with your doctor before drinking alcohol. Alcohol may make you feel faint or dizzy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Candesartan cilexetil before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Candesartan cilexetil. This medicine is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Candesartan cilexetil is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn or was born prematurely.

Driving and using machines

Some people may feel tired or dizzy when taking this medicine. If this happens to you, do not drive or use any tools or machines.

Candesartan cilexetil contains lactose

This medicine contains lactose which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3 How to take Candesartan cilexetil

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. It is important to keep taking this medicine every day. You can take Candesartan cilexetil with or without food. Swallow the tablet with a drink of water. Try to take the tablet at the same time each day. This will help you to remember to take it.

High blood pressure

- The recommended dose of Candesartan cilexetil is 8 mg once a day. Your doctor may increase this dose to 16 mg once a

<p>day and further up to 32 mg once a day depending on blood pressure response.</p> <ul style="list-style-type: none"> In some patients, such as those with liver problems, kidney problems or those who recently have lost body fluids, e.g., through vomiting or diarrhoea or by using water tablets, the doctor may prescribe a lower starting dose. Some black patients may have a reduced response to this type of medicine, when given as the only treatment, and these patients may need a higher dose. <p>Heart failure</p> <ul style="list-style-type: none"> The recommended starting dose of Candesartan cilexetil is 4 mg once a day. Your doctor may increase your dose by doubling the dose at intervals of at least 2 weeks up to 32 mg once a day. Candesartan cilexetil can be taken together with other medicines for heart failure, and your doctor will decide which treatment is suitable for you. <p>Use in children and adolescents with high blood pressure: Children 6 to <18 years of age: The recommended starting dose is 4 mg once daily. For patients weighing < 50 kg: In some patients whose blood pressure is not adequately controlled, your doctor may decide the dose needs to be increased to a maximum of 8 mg once daily. For patients weighing ≥ 50 kg: In some patients whose blood pressure is not adequately controlled, your doctor may decide the dose needs to be increased to 8 mg once daily and to 16 mg once daily. The tablet can be divided into equal doses.</p> <p>If you take more Candesartan cilexetil than you should If you take more Candesartan cilexetil than prescribed by your doctor, contact a doctor or pharmacist immediately for advice.</p> <p>If you forget to take Candesartan cilexetil Do not take a double dose to make up for a forgotten tablet. Just take the next dose as normal.</p> <p>If you stop taking Candesartan cilexetil If you stop taking Candesartan cilexetil, your blood pressure may increase again. Therefore do not stop taking this medicine without first talking to your doctor.</p> <p>If you have any further questions on the use of this medicine, ask your doctor or pharmacist.</p>	<p>Not known (frequency cannot be estimated from the available data)</p> <ul style="list-style-type: none"> Diarrhoea. <p>Additional side effects in children and adolescents In children treated for high blood pressure, side effects appear to be similar to those seen in adults, but they happen more often. Sore throat is a very common side effect in children but not reported in adults and runny nose, fever and increased heart rate are common in children but not reported in adults.</p> <p>Reporting of side effects If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.</p>
<p>4 Possible side effects</p> <p>Like all medicines, this medicine can cause side effects, although not everybody gets them. It is important that you are aware of what these side effects may be.</p> <p>Stop taking Candesartan cilexetil and seek medical help immediately if you have any of the following allergic reactions:</p> <ul style="list-style-type: none"> difficulties in breathing, with or without swelling of the face, lips, tongue and/or throat. swelling of the face, lips, tongue and/or throat, which may cause difficulties in swallowing. severe itching of the skin (with raised lumps). <p>This medicine may cause a reduction in number of white blood cells. Your resistance to infection may be decreased and you may notice tiredness, an infection or a fever. If this happens contact your doctor. Your doctor may occasionally do blood tests to check whether Candesartan cilexetil has had any effect on your blood (agranulocytosis).</p> <p>Other possible side effects include: Common (may affect up to 1 in 10 people)</p> <ul style="list-style-type: none"> Feeling dizzy/spinning sensation Headache Respiratory infection Low blood pressure. This may make you feel faint or dizzy Changes in blood test results: <ul style="list-style-type: none"> An increased amount of potassium in your blood, especially if you already have kidney problems or heart failure. If this is severe you may notice tiredness, weakness, irregular heart beat or pins and needles Effects on how your kidneys work, especially if you already have kidney problems or heart failure. In very rare cases, kidney failure may occur <p>Very rare (may affect up to 1 in 10,000 people)</p> <ul style="list-style-type: none"> Swelling of the face, lips, tongue and/or throat A reduction in your red or white blood cells. You may notice tiredness, an infection or a fever Skin rash, lumpy rash (hives) Itching Cough Back pain, pain in joints and muscles Changes in how your liver is working, including inflammation of the liver (hepatitis). You may notice tiredness, yellowing of your skin and the whites of your eyes and flu like symptoms Intestinal angioedema: a swelling in the gut presenting with symptoms like abdominal pain, nausea, vomiting and diarrhoea Nausea Changes in blood test results: <ul style="list-style-type: none"> A reduced amount of sodium in your blood. If this is severe then you may notice weakness, lack of energy, or muscle cramps 	<p>5 How to store Candesartan cilexetil</p> <p>Keep this medicine out of the sight and reach of children.</p> <p>Do not use this medicine after the expiry date which is stated on the blister foil and carton after {EXP}. The expiry date refers to the last day of that month.</p> <p>This medicine does not require any special storage conditions.</p> <p>Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.</p> <p>6 Contents of the pack and other information</p> <p>What Candesartan cilexetil contains</p> <ul style="list-style-type: none"> The active substance is Candesartan cilexetil. Each Candesartan cilexetil 8 mg Tablet contains 8 mg of candesartan cilexetil. Each Candesartan cilexetil 16 mg Tablet contains 16 mg of candesartan cilexetil. Each Candesartan cilexetil 32 mg Tablet contains 32 mg of candesartan cilexetil. The other ingredients (excipients) are pregelatinized maize starch, povidone K-30, carmellose calcium, microcrystalline cellulose (E460), lactose monohydrate, magnesium stearate, poloxamer 188 and red iron oxide (E172). <p>What Candesartan cilexetil looks like and contents of the pack</p> <ul style="list-style-type: none"> Candesartan cilexetil 8 mg Tablet is a pink, capsule shaped tablet, 7.7 mm in length and 3.5 mm in width, scored on both sides. One side of the tablet is debossed with “C 8” and the other side is debossed with “8 C”. The tablet can be divided into equal doses. Candesartan cilexetil 16 mg Tablet is a pink, capsule shaped tablet, 9.7 mm in length and 4.3 mm in width, scored on one side. One side of the tablet is debossed with the number “16”. The other side of the tablet is debossed with “C C”. The tablet can be divided into equal doses. Candesartan cilexetil 32 mg Tablet is a pink, capsule shaped tablet, 12.2 mm in length and 5.4 mm in width, scored on one side. One side of the tablet is debossed with the number “32”. The other side of the tablet is debossed with “C C”. The tablet can be divided into equal doses. <p>Candesartan cilexetil tablets are available in OPA/Aluminium/PVC – PVC/PVAC/Aluminium/OPA blisters in pack sizes of 7, 14, 15, 20, 28, 30, 50, 50 x 1 unit dose blisters (hospital pack), 56, 60, 84, 90, 98, 100, 250 & 300 tablets.</p> <p>Not all pack sizes may be marketed.</p> <p>Marketing Authorisation Holder Teva UK Limited, Ridings Point, Whistler Drive, Castleford, WF10 5HX, United Kingdom</p> <p>Manufacturer: PLIVA Hrvatska d.o.o. (PLIVA Croatia Ltd.), Prilaz baruna Filipovića 25, 10000 Zagreb, Croatia</p> <p>This leaflet was last revised in December 2024.</p> <p>PL 00289/1176 PL 00289/1177 PL 00289/1178</p>