

Vildagliptin/Metformin
50 mg/850 mg and 50 mg/1000 mg
Film-coated Tablets

vildagliptin/metformin hydrochloride

Package leaflet: Information for the user

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Vildagliptin/Metformin is and what it is used for
2. What you need to know before you take Vildagliptin/Metformin
3. How to take Vildagliptin/Metformin
4. Possible side effects
5. How to store Vildagliptin/Metformin
6. Contents of the pack and other information

1 What Vildagliptin/Metformin is and what it is used for

The active substances of Vildagliptin/Metformin, vildagliptin and metformin, belong to a group of medicines called “oral antidiabetics”.

Vildagliptin/Metformin is used to treat adult patients with type 2 diabetes. This type of diabetes is also known as non-insulin-dependent diabetes mellitus. Vildagliptin/Metformin is used when diabetes cannot be controlled by diet and exercise alone and/or with other medicines used to treat diabetes (insulin or sulphonylureas).

Type 2 diabetes develops if the body does not make enough insulin or if the insulin that the body makes does not work as well as it should. It can also develop if the body produces too much glucagon.

Both insulin and glucagon are made in the pancreas. Insulin helps to lower the level of sugar in the blood, especially after meals. Glucagon triggers the liver to make sugar, causing the blood sugar level to rise.

How Vildagliptin/Metformin works

Both active substances, vildagliptin and metformin, help to control the level of sugar in the blood. The substance vildagliptin works by making the pancreas produce more insulin and less glucagon. The substance metformin works by helping the body to make better use of insulin. This medicine has been shown to reduce blood sugar, which may help to prevent complications from your diabetes.

2 What you need to know before you take Vildagliptin/Metformin

Do not take Vildagliptin/Metformin if you:

- are allergic to vildagliptin, metformin or any of the other ingredients of this medicine (listed in section 6). If you think you may be allergic to any of these, talk to your doctor before taking Vildagliptin/Metformin.
- have uncontrolled diabetes, with, for example severe hyperglycaemia (high blood glucose), nausea, vomiting, diarrhoea, rapid weight loss, lactic acidosis (see “Risk of lactic acidosis” below) or ketoacidosis. Ketoacidosis is a condition in which substances called ketone bodies accumulate in the blood and which can lead to diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, sleepiness or your breath developing an unusual fruity smell.
- have recently had a heart attack or if you have heart failure or serious problems with your blood circulation or difficulties in breathing which could be a sign of heart problems
- have severely reduced kidney function
- have a severe infection or are seriously dehydrated (have lost a lot of water from your body)
- are going to have a contrast x-ray (a specific type of x-ray involving an injectable dye). Please also see information about this in section “Warnings and precautions”
- have liver problems
- drink alcohol excessively (whether every day or only from time to time)
- are breast-feeding (see also “Pregnancy and breast-feeding”).

Warnings and precautions

Talk to your doctor or pharmacist before taking
Vildagliptin/Metformin

Risk of lactic acidosis

Vildagliptin/Metformin may cause a very rare, but very serious side effect called lactic acidosis, particularly if your kidneys are not working properly. The risk of developing lactic acidosis is also increased with uncontrolled diabetes, serious infections, prolonged fasting or alcohol intake, dehydration (see further information below), liver problems and any medical conditions in which a part of the body has a reduced supply of oxygen (such as acute severe heart disease).
If any of the above apply to you, talk to your doctor for further instructions.

Stop taking Vildagliptin/Metformin for a short time if you have a condition that may be associated with dehydration (significant loss of body fluids) such as severe vomiting, diarrhoea, fever, exposure to heat or if you drink less fluid than normal. Talk to your doctor for further instructions.

Stop taking Vildagliptin/Metformin and contact a doctor or the nearest hospital immediately if you experience some of the symptoms of lactic acidosis, as this condition may lead to coma. Symptoms of lactic acidosis include:

- vomiting
- stomach ache (abdominal pain)
- muscle cramps
- a general feeling of not being well with severe tiredness
- difficulty in breathing
- reduced body temperature and heartbeat.

Lactic acidosis is a medical emergency and must be treated in a hospital.

Talk to your doctor promptly for further instructions if:

- You are known to suffer from a genetically inherited disease affecting mitochondria (the energy-producing components within cells) such as MELAS syndrome (Mitochondrial Encephalopathy, myopathy, Lactic acidosis and Stroke-like episodes) or Maternal inherited diabetes and deafness (MIDD).
- You have any of these symptoms after starting metformin: seizure, declined cognitive abilities, difficulty with body movements, symptoms indicating nerve damage (e.g. pain or numbness), migraine and deafness.

Vildagliptin/Metformin is not a substitute for insulin. Therefore, you should not receive Vildagliptin/Metformin for the treatment of type 1 diabetes.

Talk to your doctor, pharmacist or nurse before taking Vildagliptin/Metformin if you have or have had a disease of the pancreas.

Talk to your doctor, pharmacist or nurse before taking Vildagliptin/Metformin if you are taking an anti-diabetic medicine known as a sulphonylurea. Your doctor may want to reduce your dose of the sulphonylurea when you take it together with Vildagliptin/Metformin in order to avoid low blood glucose (hypoglycaemia).

If you have previously taken vildagliptin but had to stop taking it because of liver disease, you should not take this medicine.

Diabetic skin lesions are a common complication of diabetes. You are advised to follow the recommendations for skin and foot care that you are given by your doctor or nurse. You are also advised to pay particular attention to new onset of blisters or ulcers while taking Vildagliptin/Metformin. Should these occur, you should promptly consult your doctor.

If you need to have major surgery you must stop taking Vildagliptin/Metformin during and for some time after the procedure. Your doctor will decide when you must stop and when to restart your treatment with Vildagliptin/Metformin.

A test to determine your liver function will be performed before the start of Vildagliptin/Metformin treatment, at three-month intervals for the first year and periodically thereafter. This is so that signs of increased liver enzymes can be detected as early as possible.

During treatment with Vildagliptin/Metformin, your doctor will check your kidney function at least once a year or more frequently if you are elderly and/or have worsening renal function.

Your doctor will test your blood and urine for sugar regularly.

Children and adolescents

The use of Vildagliptin/Metformin in children and adolescents up to 18 years of age is not recommended.

Other medicines and Vildagliptin/Metformin

If you need to have an injection of a contrast medium that contains iodine into your bloodstream, for example in the context of an X-ray or scan, you must stop taking Vildagliptin/Metformin before or at the time of the injection. Your doctor will decide when you must stop and when to restart your treatment with Vildagliptin/Metformin.

Tell your doctor if you are taking, have recently taken or might take any other medicines. You may need more frequent blood glucose and kidney function tests, or your doctor may need to adjust the dosage of Vildagliptin/Metformin. It is especially important to mention the following:

- glucocorticoids generally used to treat inflammation
- beta-2 agonists generally used to treat respiratory disorders
- other medicines used to treat diabetes
- medicines which increase urine production (diuretics)
- medicines used to treat pain and inflammation (NSAID and COX-2-inhibitors, such as ibuprofen and celecoxib)
- certain medicines for the treatment of high blood pressure (ACE inhibitors and angiotensin II receptor antagonists)
- certain medicines affecting the thyroid
- certain medicines affecting the nervous system
- certain medicines used to treat angina (e.g. ranolazine)
- certain medicines used to treat HIV infection (e.g. dolutegravir)
- certain medicines used to treat a specific type of thyroid cancer (medullary thyroid cancer) (e.g. vandetanib)
- certain medicines used to treat heartburn and peptic ulcers (e.g. cimetidine).

Vildagliptin/Metformin with alcohol

Avoid excessive alcohol intake while taking Vildagliptin/Metformin since this may increase the

risk of lactic acidosis (please see section “Warnings and precautions”).

Pregnancy and breast-feeding

- If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildagliptin/Metformin during pregnancy.
- Do not use Vildagliptin/Metformin if you are pregnant or breast-feeding (see also “Do not take Vildagliptin/Metformin”).

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

If you feel dizzy while taking Vildagliptin/Metformin, do not drive or use any tools or machines.

3 How to take Vildagliptin/Metformin

The amount of Vildagliptin/Metformin that people have to take varies depending on their condition. Your doctor will tell you exactly the dose of Vildagliptin/Metformin to take.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is one film-coated tablet of either 50 mg/850 mg or 50 mg/1000 mg taken twice a day.

If you have reduced kidney function, your doctor may prescribe a lower dose. Also if you are taking an anti-diabetic medicine known as a sulphonylurea your doctor may prescribe a lower dose.

Your doctor may prescribe this medicine alone or with certain other medicines that lower the level of sugar in your blood.

When and how to take Vildagliptin/Metformin

- Swallow the tablets whole with a glass of water.
- Take one tablet in the morning and the other in the evening with or just after food. Taking the tablet just after food will lower the risk of an upset stomach.

The I shaped score line is only there to help you break the tablet if you have difficulty swallowing it whole.

Continue to follow any advice about diet that your doctor has given you. In particular, if you are following a diabetic weight control diet, continue with this while you are taking Vildagliptin/Metformin.

If you take more Vildagliptin/Metformin than you should

If you take too many Vildagliptin/Metformin tablets, or if someone else takes your tablets, **talk to a doctor or pharmacist immediately**. Medical attention may be necessary. If you have to go to a doctor or hospital, take the pack and this leaflet with you.

If you forget to take Vildagliptin/Metformin

If you forget to take a tablet, take it with your next meal unless you are due to take one then anyway. Do not take a double dose (two tablets at once) to make up for a forgotten tablet.

If you stop taking Vildagliptin/Metformin

Continue to take this medicine as long as your doctor prescribes it so that it can continue to control your blood sugar. Do not stop taking Vildagliptin/Metformin unless your doctor tells you to. If you have any questions about how long to take this medicine, talk to your doctor.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You should **stop taking Vildagliptin/Metformin and see your doctor immediately** if you experience the following side effects:

- **Lactic acidosis (very rare: may affect up to 1 in 10 000 people):** Vildagliptin/Metformin may cause a very rare, but very serious side effect called lactic acidosis (see section “Warnings and precautions”). If this happens you must **stop taking Vildagliptin/Metformin and contact a doctor or the nearest hospital immediately**, as lactic acidosis may lead to coma. Signs of a high level of lactic acid in the blood include drowsiness or dizziness, severe nausea or vomiting, abdominal pain, irregular heart beat or deep, rapid breathing.
- **Angioedema (rare: may affect up to 1 in 1 000 people):** Symptoms include swollen face, tongue or throat, difficulty swallowing, difficulty breathing, sudden onset of rash or hives, which may indicate a reaction called “angioedema”.
- **Liver disease (hepatitis) (uncommon: may affect up to 1 in 100 people):** Symptoms include yellow skin and eyes, nausea, loss of appetite or dark-coloured urine, which may indicate liver disease (hepatitis).
- **Inflammation of the pancreas (pancreatitis) (uncommon: may affect up to 1 in 100 people):** Symptoms include severe and persistent pain in the abdomen (stomach area), which might reach through to your back, as well as nausea and vomiting.

Other side effects

Some patients have experienced the following side effects while taking Vildagliptin/Metformin:

- **Common (may affect up to 1 in 10 people):** sore throat, runny nose, fever, itchy rash, excessive sweating, joint pain, dizziness, headache, trembling that cannot be controlled, weakness, constipation, nausea (feeling sick), vomiting, diarrhoea, flatulence, heartburn, pain in and around the stomach (abdominal pain).
- **Uncommon (may affect up to 1 in 100 people):** tiredness, metallic taste, low blood glucose, loss of appetite, swollen hands, ankles or feet (oedema), chills, muscle pain, redness of the skin, urticaria.
- **Very rare (may affect up to 1 in 10 000 people):** decreased vitamin B12 levels (paleness, tiredness, mental symptoms such as confusion or memory disturbances).

Since this product has been marketed, the following side effects have also been reported:

- **Not known: (frequency cannot be estimated from the available data):** localised peeling of skin or blisters, blood vessel inflammation (vasculitis) which may result in skin rash or pointed, flat, red, round spots under the skin's surface or bruising.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Vildagliptin/Metformin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister foil after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C.
Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Vildagliptin/Metformin contains

- The active substances are vildagliptin and metformin hydrochloride. Each Vildagliptin/Metformin 50 mg/850 mg film-coated tablet contains 50 mg vildagliptin and 850 mg metformin hydrochloride (corresponding to 660 mg of metformin). Each Vildagliptin/Metformin 50 mg/1000 mg film-coated tablet contains 50 mg vildagliptin and 1000 mg metformin hydrochloride (corresponding to 780 mg of metformin).
- The other ingredients are:
Tablet core: Hydroxypropylcellulose, microcrystalline cellulose, magnesium stearate.
Film-coating: Hypromellose 2910, 6 cps, titanium dioxide (E171), copovidone, polydextrose, macrogol 3350, yellow iron oxide (E172), medium-chain triglycerides.

What Vildagliptin/Metformin looks like and contents of the pack

Vildagliptin/Metformin 50 mg/850 mg film-coated tablets are yellow, oval and biconvex shaped tablets, 8.5 x 21 mm in size with an I-shaped score line on one side and debossed with “A 8” on the other side.

The I-shaped score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

Vildagliptin/Metformin 50 mg/1000 mg film-coated tablets are brownish yellow, oval and biconvex shaped tablets, 9.2 x 23 mm in size with an I-shaped score line on one side and debossed with “A 1” on the other side.

The I-shaped score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

Vildagliptin/Metformin is available in blister packs containing 10, 12, 30, 60, 100 and 180 film-coated tablets, multipacks comprising 2 cartons, each containing 30 film-coated tablets and in perforated unit dose blister packs containing 30 x 1 and 60 x 1 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder


Teva UK Limited, Ridings Point, Whistler Drive, Castleford, WF10 5HX, United Kingdom

Manufacturer

Balkanpharma-Dupnitsa AD, 3 Samokovsko Shosse Str., Dupnitsa, 2600, Bulgaria

This leaflet was last revised in 03/2025

PL 00289/2174-2175

238179_s1